

**Erratum: “Doppler Broadening and its Contribution to Compton Energy-Absorption Cross Sections: An Analysis of the Compton Component in Terms of Mass-Energy Absorption Coefficient”**  
[J. Phys. Chem. Ref. Data 31, 769 (2002)]

**D. V. Rao,<sup>a)</sup> T. Takeda, and Y. Itai**

*Institute of Clinical Medicine, University of Tsukuba, Ibaraki 305-8575, Japan*

**T. Akatsuka**

*Faculty of Engineering, Yamagata University, Yonezawa-shi, Yamagata 992-8510, Japan*

**R. Cesareo and A. Brunetti**

*Instituto de Matematica e Fisica, Universita di Sassari, Via Vienna, 2-1-07100 Sassari, Italy*

**G. E. Gigante**

*Dipartimento di Fisica, Universita di Roma “La Sapienza,” Ple. A Moro2, 00185 Roma, Italy*

(Published 28 May 2003)

The captions for Fig. 1 and Table 3 and the y-axis label for Fig. 3 were incorrect. The corrections are as follows.

Page 773: Fig. 1. A comparison of the present values with the standard theoretical estimates: (1) F. Biggs *et al.*<sup>9</sup> [At. Data Nucl. Tables **16**, 201 (1975)]. (2) E. Storm and H.

I. Israel<sup>10</sup> [Nucl. Data Tables **A7**, 565 (1970)]. (3) present values (a) hydrogen, (b) oxygen, (c) sulfur, and (4) calcium.

Page 775: Fig. 3. Label for the y axis should be  $\Delta G(\omega_1)$  (keV).

Page 779: Table 3. Compton component of the mass-energy absorption coefficient ( $\text{cm}^2 \cdot \text{g}^{-1}$ ) in the region from 1 keV to 1 MeV.

---

<sup>a)</sup>Electronic mail: dvrao@ssmain.uniss.it